

This note, that [was found here](#), as a part of [a collection](#) is written (completely with human hands) by [Rupadarshi Ray](#), was created on March 4, 2026 11:04:54 PM, was last modified on April 7, 2026 3:22:38 AM, and was exported on September 7, 2026 3:09:48 PM.

2-generated Kleinian groups

✓ <https://ariymarkowitz.github.io/H3-Visualiser/?d=5&a=1.922423900484707+-1.989954347180768+10+0+1+0+1+0&b=4+0+6+0+-0.8940552095531961+0.7324313470166844+10+0&showa=1&showb=1>

Current note has 0 direct children and 0 total descendants.

- stem
 - Objects stemming from $PSL(2)(\mathbb{C})$
 - Kleinian groups $\Gamma <_d PSL(2)(\mathbb{C}) \leftrightarrow$ Hyperbolic 3-manifolds/orbifolds $\Gamma \backslash \mathbb{R}H^3$
 - 2-generated Kleinian groups

And it has 7 siblings.

- stem
 - Objects stemming from $PSL(2)(\mathbb{C})$
 - Kleinian groups $\Gamma <_d PSL(2)(\mathbb{C}) \leftrightarrow$ Hyperbolic 3-manifolds/orbifolds $\Gamma \backslash \mathbb{R}H^3$
 - Finitely generated, free Kleinian groups
 - 2-generated Kleinian groups
 - Finitely generated Kleinian groups $\leftrightarrow$ topologically tame oriented hyperbolic 3-manifolds/finite degree orbifold covers of
 - Building hyperbolic 3-manifolds by gluing polyhedra
 - Kleinian group with $\Lambda(\Gamma) \neq \partial \mathbb{R}H^3$
 - Thick-thin decomposition of hyperbolic 3-manifolds
 - Kleinian ultra-Schottky groups $\leftrightarrow$ convex cocompact hyperbolic interior of 3-handlebodies