

 Info

This note, that [was found here](#), as a part of [a collection](#) is written (completely with human hands) by [Rupadarshi Ray](#), was created on December 16, 2025 11:02:51 PM, was last modified on September 6, 2026 3:21:40 PM, and was exported on September 7, 2026 3:24:51 PM.

Closed map between topological spaces

 **Definition.** Closed maps, maps with continuity of preimage

A map $f : X \rightarrow Y$ between topological spaces

- is **closed** if for every closed set $M \subseteq X$ the image $f(M) \subseteq Y$ is closed
- has **continuity of the preimage** if for all $y \in Y$ and open nb $U \subseteq X$ of $f^{-1}(y)$, there exists a open nb $V \subseteq Y$ of y such that

$$f^{-1}(V) \subseteq U$$

Inclusion maps $\iota : A \hookrightarrow X$ are closed $\iff$ for every closed subset $S \subseteq A$

$$\iota(S) = A \cap S$$

is closed $\iff A \subseteq X$ is closed.

 Projections

$$X \times Y \rightarrow X$$

are not closed in general.

Consider

$$\pi_1 : \mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$$

Then

$$\underbrace{V(xy - 1)}_{\text{closed}} \mapsto \underbrace{(-\infty, 0) \cup (0, \infty)}_{\text{not closed}}$$

 Let Y be a topological space. Then Y is compact $\iff$ for every topological space X the projection

$$\pi_1 : X \times Y \rightarrow X$$

is **closed**.

 Local homeomorphisms, even covering maps, are not closed in general.

Consider the covering

$$\begin{aligned} \mathbb{R} &\rightarrow S^1 \\ x &\mapsto \exp(2\pi i x) \end{aligned}$$

We know

$$f^{-1}(1) = \mathbb{Z}$$

and we have an evenly covered nb of $\mathbb{Z}$ mapping to 1

$$\bigsqcup_{m \in \mathbb{Z}} \left(\left(-\frac{1}{2}, \frac{1}{2} \right) + m \right) = f^{-1}(1)$$

Also note

$$\frac{1}{n+2} \xrightarrow{n \geq 1, n \rightarrow \infty} 0 \implies \exp\left(\frac{2\pi i}{n+2}\right) \xrightarrow{n \geq 1, n \rightarrow \infty} 1$$

Then, as this is an infinitely sheeted cover, we pick a preimage of each element of the converging sequence in *different* sheets

$$\frac{1}{|m|+2} + m \in \left(-\frac{1}{2}, \frac{1}{2} \right) + m$$

implying

$$\left\{ \frac{1}{|m|+2} + m \mid m \in \mathbb{Z} \right\}$$

is a discrete, closed set.

$$\underbrace{\left\{ \frac{1}{|m|+2} + m \mid m \in \mathbb{Z} \right\}}_{\text{closed} \subseteq \mathbb{R}} \mapsto \underbrace{\left\{ \exp\left(\frac{2\pi i}{n+2}\right) \mid n \in \mathbb{N} \right\}}_{\text{not closed} \subseteq S^1}$$

closed $\iff$ continuity of preimage

 Let $f : X \rightarrow Y$ be map between topological spaces. Then f is **closed** $\iff$ f has **continuity of the preimage**.

 Let f be closed.

- For $y \in Y$ and U be a open nb of $f^{-1}(y)$. Then

$$f(X \setminus U)$$

is closed and does not contain y .

- Hence, there is a nb V of y such that

$$V \cap f(X \setminus U) = \emptyset \iff f^{-1}(V) \subseteq U$$

