

Info

This note, that [was found here](#), as a part of [a collection](#) is written (completely with human hands) by [Rupadarshi Ray](#), was created on January 14, 2025 1:57:29 PM, was last modified on September 7, 2026 1:11:14 PM, and was exported on September 7, 2026 3:09:57 PM.

$$\mathcal{O}(U)$$

Let $U \subseteq \mathbb{C}$ be open and

$$\mathcal{O}(U)$$

be the $\mathbb{C}$ -algebra of holomorphic functions.

$$\mathcal{O}(U) \subseteq \mathcal{C}(U, \mathbb{C})$$

From

Definition. $\mathcal{C}(U, X)$

Let (X, d) be a complete metric space, $U \subseteq \mathbb{R}^n$ be a open subset and

$$\text{EndTop}(U, X)$$

be the space of all continuous functions $U \rightarrow X$.

- For $K \subseteq U$ compact we have the distance

$$\rho_K(f, g) := \sup \{d(f(z), g(z)) | z \in K\}$$

for all $f, g \in \mathcal{C}(G, X)$.

- Then for $U = \bigcup_{n \in \mathbb{N}} K_n$ where K_n is compact and $K_n \subseteq \text{int}(K_{n+1})$ we define the metric

$$\rho(f, g) := \sum_{n \in \mathbb{N}} \frac{1}{2^n} \frac{\rho_{K_n}(f, g)}{1 + \rho_{K_n}(f, g)}$$

on $\text{EndTop}(U, X)$.

The (metrizable) topological space $\text{EndTop}(U, X)$ with the topology induced by the metric ρ is denoted $\mathcal{C}(U, X)$.

we have the compact-open topology on

$$\mathcal{O}(U) \subseteq \mathcal{C}(U, \mathbb{C})$$

If $\{f_n\}$ is a sequence in $\mathcal{O}(U)$ and $f \in \mathcal{C}(U, \mathbb{C})$ such that $f_n \xrightarrow{n \rightarrow \infty} f$ then $f \in \mathcal{O}(U)$ and all derivatives

$$f_n^{(k)} \xrightarrow{n \rightarrow \infty} f^{(k)}$$

for $k \geq 1$. Thus

$$\mathcal{D} : \mathcal{O}(U) \rightarrow \mathcal{O}(U)$$

is a continuous linear map.

connected $U \iff \mathcal{O}(U)$ is an integral domain

An open set $U \subseteq \mathbb{C}$ is connected $\iff \mathcal{O}(U)$ is an [integral domain](#).

Let

$$f, g : U \rightarrow \mathbb{C}$$

be holomorphic on such that

$$fg \equiv 0 \text{ on } U$$

and $f(z_0) \neq 0$ for some $z_0 \in U$. Then $f \neq 0$ on a disk around z_0 , and thus $g = 0$ on that disk. Therefore by

(Identity theorem for holomorphic functions on subsets of $\mathbb{C}$) Let

$$f : \Omega(\text{open}) \subseteq \mathbb{C} \rightarrow \mathbb{C}$$

be holomorphic. And L be the set of limit points of $f^{-1}(0)$. Then L is **both open and closed in Ω** . Therefore if Ω is connected, either $L = \Omega \iff f \equiv 0$ on Ω or $f^{-1}(0)$ has no limit point in Ω , that is if $f^{-1}(0)$ has a limit point in Ω then $f \equiv 0$.

$$g \equiv 0 \text{ on } \Omega$$

- If U is not connected, define locally constant functions such that are not zero but one of f, g is zero on each component, thus $fg = 0$.

GCDs exist in $\mathcal{O}(U)$

Let

$$\mathcal{F} \subseteq \mathcal{O}(U)$$

be a non-empty subfamily but $\mathcal{F} \neq \{0\}$. We apply

(Holomorphic function on open subsets of $\mathbb{C}P^1$ with given zeros and their multiplicities) Let Ω be a proper open subset of $\mathbb{C}P^1$ and $A \subseteq \Omega$ with no limit point in Ω . For each $a \in A$, let $m_a \in \mathbb{Z}_{>0}$ be a positive integer. Then there exists a holomorphic function

$f : \Omega \rightarrow \mathbb{C}$

such that $A = f^{-1}(0)$ and for each $a \in A$ the multiplicity of f at a is m_a .

Ash and Novinger - Complex Variables, p.144

to obtain $g \in \mathcal{O}(U)$ such that

$$g^{-1}(0) = B := \bigcap \{f^{-1}(0) \mid f \in \mathcal{F}\}$$

and for each $b \in B$ the multiplicity

$$m(g,b) = \min\{m(f,b) \mid f \in \mathcal{F}\}$$

Then $f \in \mathcal{F}$ implies

$$\frac{g}{f}$$

has a removable singularity at every zero of f , thus

$$g \mid f$$

Furthermore, if $h \in \mathcal{O}(U)$ and $h \mid f$ for each $f \in \mathcal{F}$ then $h^{-1}(0) \subseteq B$ and for each $b \in B$

$$m(h,b) \leq \min\{m(f,b) \mid f \in \mathcal{F}\} = m(g,b)$$

Thus

$$\frac{h}{g}$$

extends holomorphically to U and thus $h \mid g$. Thus g is the greatest common divisor of $\mathcal{F}$.

finitely generated ideals are principal
 never a PID, never Noetherian

[1]

functor produces an equivalence of categories

Definition.

$\mathcal{O} : \text{Open}(\mathbb{C})^{\text{op}} \rightarrow \text{Alg}_{\mathbb{C}}$

Proposition: Let U_1, U_2 be open subsets of $\mathbb{C}$.

- Let $f : U_1 \rightarrow U_2$ be a holomorphic map. Then the pullback

$$f^* : \mathcal{O}(U_2) \rightarrow \mathcal{O}(U_1)$$

is a unital $\mathbb{C}$ -algebra homomorphism. The map f is a biholomorphism $\iff f^*$ is an isomorphism.
- Let $\Phi : \mathcal{O}(U_2) \rightarrow \mathcal{O}(U_1)$ be a unital $\mathbb{C}$ -algebra homomorphism. Then for $f := \Phi(\text{Id}_{U_2})$ we have

$$f : U_1 \rightarrow U_2$$

and $\Phi = f^*$.
- Therefore, $\mathcal{O}(U_1)$ and $\mathcal{O}(U_2)$ are isomorphic $\iff U_1, U_2$ are biholomorphic.

[2]

For $z \in \mathbb{C} \setminus U_2$ we have $\text{Id}_{U_2} - z \in \mathcal{O}(U_2)^\times$. Then $\Phi(\text{Id}_{U_2} - z) \in \mathcal{O}(U_1)^\times$.

- Therefore, for $f(x) := \Phi(\text{Id}_{U_2})(x)$ we have

$$f - z \in \mathcal{O}(U_1)^\times$$

implying $z \notin f(D)$.
- Thus,

$$f : U_1 \rightarrow U_2$$
- Let $g \in \mathcal{O}(U_2)$. Fix $z \in U_1$. Then

$$g - g(f(z))$$

has a zero at $f(z)$. There exists $h \in \mathcal{O}(U_2)$ such that

$$g - g(f(z)) = (\text{Id}_{U_2} - f(z))h$$
- Thus,

$$\Phi(g - g(f(z))) = (f - f(z))\Phi(h)$$

vanishes at z . Hence,

$$\Phi(g)(z) = g(f(z)) = (f^*g)(z)$$

Definition.

$\mathcal{O} : \text{Open}(\mathbb{C})^{\text{op}} \rightarrow \text{Alg}_{\mathbb{C}}$

is an equivalence of categories onto some full subcategory of $\text{Alg}_{\mathbb{C}}$.

maxSpec $\mathcal{O}(U) \cong_{\text{Set}} U$?

continuous dual

$$(\mathcal{O}(U), \mathfrak{T}_{\text{cpt}})^* \cong? \mathcal{O}_0(\mathbb{C}P^1 \setminus U)$$

- subobjects of and functions on $\mathbb{R}^n, S^n, \mathbb{C}^n$ and its quotients $\Gamma \backslash \mathbb{R}^n, \Gamma \backslash \mathbb{C}^n$
 - Holomorphic functions on spaces over $\mathbb{C}$ of dimension 1
 - $\mathcal{O}(U)$
 - Pre-compact subsets of $\mathcal{O}(U)$
 - Sequences of holomorphic functions

And it has 19 siblings.

- stem
 - subobjects of and functions on $\mathbb{R}^n, S^n, \mathbb{C}^n$ and its quotients $\Gamma \backslash \mathbb{R}^n, \Gamma \backslash \mathbb{C}^n$
 - Holomorphic functions on spaces over $\mathbb{C}$ of dimension 1
 - Approximation of holomorphic functions on $\mathbb{C}$ by rational functions
 - Embedding unit disk D into $\mathbb{C}^3$
 - Factorization of holomorphic functions on $\mathbb{C}$
 - Global holomorphic functions
 - Equivalent descriptions of holomorphic functions
 - Meromorphic functions on the complex plane
 - Holomorphic functions meromorphic at infinity
 - Modular forms
 - Reconstructing meromorphic functions from stalks
 - Extending holomorphic functions by reflections
 - Rotational symmetrization of holomorphic functions
 - Sheaf of holomorphic functions on $\mathbb{C}$
 - $\mathcal{O}(U)$
 - $\mathcal{O}(\mathbb{C})$
 - $\mathcal{O}(D)$
 - $\mathcal{O}^p(H^2_U)$
 - $\mathcal{O} \cap L^p$
 - $\mathcal{O}(S^1)$
 - Zeros and singularities of holomorphic functions

1. [Wayback Machine](#) ↩
2. [abstract algebra - Are the rings of holomorphic functions over a domain \$D \subset \mathbb{C}\$ isomorphic for different \$D\$? - Mathematics Stack Exchange](#) ↩