

This note, that [was found here](#), as a part of [a collection](#) is written (completely with human hands) by [Rupadarshi Ray](#), was created on February 4, 2026 12:10:51 AM, was last modified on February 15, 2026 11:45:12 PM, and was exported on September 7, 2026 3:10:02 PM.

# Rigidity of lattices using quasi-conformal mappings

Current note has 0 direct children and 0 total descendants.

- [stem](#)
  - [O+ R1 n](#)
    - [Discrete subgroups of  \$O^+\(1, n\)\$  ↔ hyperbolic  \$n\$ -manifolds/orbifolds](#)
      - [Rigidity of  \$\mathbb{R}\$ -hyperbolic manifolds](#)
        - [Rigidity of lattices using quasi-conformal mappings](#)

And it has 3 siblings.

- [stem](#)
  - [O+ R1 n](#)
    - [Discrete subgroups of  \$O^+\(1, n\)\$  ↔ hyperbolic  \$n\$ -manifolds/orbifolds](#)
      - [Rigidity of  \$\mathbb{R}\$ -hyperbolic manifolds](#)
        - [Using ergodic current on  \$\partial^2\mathbb{R}^n\$](#)
        - [Rigidity using measure homology](#)
        - [Rigidity of lattices using quasi-conformal mappings](#)