

This note, that [was found here](#), as a part of [a collection](#) is written (completely with human hands) by [Rupadarshi Ray](#), was created on December 23, 2024 8:02:19 PM, was last modified on February 18, 2026 1:41:04 AM, and was exported on September 7, 2026 3:25:40 PM.

Measure preserving endomorphisms

title: Measurable endomorphisms	
-----	-----

non-singular	$\mu(A) = 0 \iff \mu(T^{-1}A)=0$
conservative	if $\mu(A)>0$ then there exists $n \in \mathbb{N}$ such that $\mu(T^{-n}A \cap A)>0$ $\iff$ every wandering set has measure zero
dissipative	if it is not conservative
totally dissipative	for every $A \in \mathcal{M}$ $\mu(\bigcup_{n \in \mathbb{N}} T^{-n}A) = \mu(A)$ $\iff$ every wandering set has measure zero
incompressible	if $A \subseteq T^{-1}A$ then $\mu(T^{-1}A) = \mu(A)$

Definition. Measure preserving endomorphisms

Let $(X, \mathfrak{M}, \mu)$ be a $[0, \infty]$ -measure space and

$$T : X \rightarrow X$$

be a measurable map. Then T preserves the measure μ that is μ is a **T -invariant measure** or T is **μ -preserving transformation** if

$$A \in \mathcal{A} \implies \mu(T^{-1}(A)) = \mu(A)$$

Definition. Ergodic endomorphisms

A μ -preserving transformation $T : X \rightarrow X$ on a measure space $(X, \mathcal{A}, \mu)$ is called **μ -ergodic** if

$$A \in \mathcal{A}, T^{-1}(A) = A \implies \mu(A) = 0 \text{ or } \mu(X \setminus A) = 0$$

Definition. Incompressible transformation on a measure space

A measurable function

$$T : X \rightarrow X$$

on a measure space $(X, \mathcal{A}, \mu)$ is called **incompressible** if

$$A \in \mathcal{A}, A \subseteq T^{-1}(A) \implies \mu(T^{-1}(A)) = \mu(A)$$

Definition. Strongly mixing endomorphisms

A μ -preserving transformation $T : X \rightarrow X$ on a **probability** space $(X, \mathcal{A}, \mu)$ is called **strongly mixing** if

$$A, B \in \mathcal{A} \implies \mu(A \cap T^{-n}(B)) \xrightarrow{n \rightarrow \infty} \mu(A)\mu(B)$$

ergodic	for every $A \in \mathfrak{M}$ with $\mu(A) > 0$ there is a $n \geq 1$ such that $\mu(T^n(A) \cap A) > 0$
strongly mixing	if for all $A, B \in \mathfrak{M}$ we have $\mu(A \cap T^{-n}B) \xrightarrow{n \rightarrow \infty} \mu(A)\mu(B)$
weakly mixing	$\sum_{n=0}^{N-1} \frac{1}{N} \mu(A \cap T^{-n}B) - \mu(A)\mu(B) \xrightarrow{N \rightarrow \infty} 0$

Strong mixing on a probability space $\implies$ ergodicity.

Suppose $A \in \mathcal{A}$ such that $T^{-1}(A) = A$. Then

$$\mu(E) = \mu(E \cap T^{-n}(E)) \xrightarrow{n \rightarrow \infty} \mu(E)^2$$

which means

$$\mu(E)^2 = \mu(E) \implies \mu(E) = 1 \text{ or } 0$$

Current note has 2 direct children and 2 total descendants.

- [structures based on sets](#)
 - [Measure spaces](#)
 - [Measure preserving endomorphisms](#)
 - [Interpretation of measure preserving endomorphisms](#)
 - [Recurrence in iterations of a measure preserving endomorphism](#)

And it has 7 siblings.

- [structures based on sets](#)
 - [Measure spaces](#)
 - [Algebra of subsets of a set](#)
 - [Meas](#)
 - [Functions on measure spaces](#)
 - [ℂ-measures](#)
 - [\[0, ∞\]-measure spaces](#)
 - [Measure preserving endomorphisms](#)
 - [\[0, 1\]-measure spaces](#)