

This note, that [was found here](#), as a part of [a collection](#) is written (completely with human hands) by [Rupadarshi Ray](#), was created on December 9, 2025 7:33:06 PM, was last modified on December 9, 2025 7:34:51 PM, and was exported on September 7, 2026 3:31:21 PM.

Geodesic flow on Riemannian manifolds of negative curvature

Let (M, g) be a compact Riemannian manifold of negative curvature. Then the geodesic flow on $T^u M$ is **ergodic**.

Current note has 0 direct children and 0 total descendants.

- [structures based on sets](#)
 - [manifolds](#)
 - [ℝ-manifold semi-Riemannian](#)
 - [Complete Riemannian manifolds](#)
 - [Riemannian manifolds with negative curvature](#)
 - [Geodesic flow on Riemannian manifolds of negative curvature](#)

And it has 4 siblings.

- [structures based on sets](#)
 - [manifolds](#)
 - [ℝ-manifold semi-Riemannian](#)
 - [Complete Riemannian manifolds](#)
 - [Riemannian manifolds with negative curvature](#)
 - [Geodesic flow on Riemannian manifolds of negative curvature](#)
 - [Discrete subgroup of isometry_group of a simply connected Riemannian manifold of negative curvature](#)
 - [Rigidity of marked length spectrum for Riemannian 2-manifolds of negative curvature](#)
 - [Equivalence of marked length spectrum, geodesic flow and boundary of universal cover of Riemannian manifolds with negative curvature](#)