

## Info

This note [found here](#)  
as a part of [a collection](#)  
is written (completely with human hands) by [Rupadarshi Ray](#),  
created on June 18, 2024 1:12:04 AM,  
and was last modified on June 12, 2026 7:35:21 PM.

# Modular forms

## Definition. Weakly modular forms of level $SL_{\mathbb{Z}}(2)$ on upper-half plane

Let  $k \in \mathbb{N}$ . A meromorphic function

$$f : H \rightarrow \mathbb{C}$$

is **weakly modular** of weight  $k$  if

$$\tau \in \mathbb{H}, \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in SL_{\mathbb{Z}}(2) \implies f\left(\begin{bmatrix} a & b \\ c & d \end{bmatrix}(\tau)\right) = (c\tau + d)^k f(\tau)$$

- If this rule is satisfied for  $S = \begin{bmatrix} 1 & 1 \\ 0 & 1 \end{bmatrix}, \begin{bmatrix} 1 & -1 \\ 1 & 0 \end{bmatrix}$  then it holds for all  $SL_{\mathbb{Z}}(2)$ . This means  $f$  is weakly modular of weight  $k$  if

$$f(\tau + 1) = f(\tau) \text{ and } f\left(-\frac{1}{\tau}\right) = \tau^k f(\tau)$$

- Weakly modular functions are  $\mathbb{Z}\{1\}$ -periodic.
- Weakly modular of **weight 0** is simply  $SL_{\mathbb{Z}}(2)$ -invariance of  $f$ .
- Weakly modular of **weight 2** is simply  $SL_{\mathbb{Z}}(2)$ -invariance of

$$f(\tau)d\tau$$

## modular forms

•

$$H \rightarrow D \setminus \{0\}$$
$$\tau \mapsto e^{2\pi i \tau}$$

is a holomorphic  $\mathbb{Z}$ -periodic map

•


$$\Im \tau \rightarrow \infty \iff q = e^{2\pi i \tau} \rightarrow 0$$

- $\Im\tau \rightarrow 0 \iff |q| \rightarrow 1$
- $\partial H = \mathbb{R}$  "goes to"  $\partial D = S^1$
- We have the inverse

Definition. The complex logarithm

$$\mathbb{C} \setminus (-\infty, 0] \rightarrow \mathbb{C}$$

$$\log(z) := \int_{1 \rightarrow z} \frac{dw}{w}$$



on

$$D \setminus \mathbb{R}_{\leq 0} \rightarrow H$$

- If

$$f: H \rightarrow \mathbb{C}$$

is holomorphic/meromorphic such that  $f(z+1) = f(z)$  ( $\mathbb{Z}$ -periodic)

$$\iff$$

there is a holomorphic/meromorphic  $g: \mathbb{D}^* \rightarrow \mathbb{C}$  such that

$$f(\tau) = g(e^{2\pi i \tau})$$

given by the composition of log

$$g = f \circ \frac{\log}{2\pi i}$$

which is well-defined as

- two different branches of log differ by  $2\pi i k$  then

$$f\left(\frac{\log q + 2\pi i k}{2\pi i}\right) = f\left(\frac{\log q}{2\pi i}\right)$$

! match

- Now  $g$  has a Laurent series

$$g = \sum_{n \in \mathbb{Z}} a_n q^n \text{ on } D \setminus \{0\}$$

- which gives  $f$  has a Fourier expansion

$$f(\tau) = \sum_{n \in \mathbb{Z}} a_n e^{(2\pi i)n\tau}$$

### Definition. Holomorphicity at $\infty$ of $\mathbb{Z}$ -periodic functions on $H$

The holomorphic

$$f : H \rightarrow \mathbb{C}$$

is **holomorphic at  $\infty$**  if the corresponding

$$g : D \setminus \{0\} \rightarrow \mathbb{C}$$

extends to  $0 \in D$ . This is equivalent to existence of

$$\lim_{\text{im}(\tau) \rightarrow \infty} f(\tau)$$

or even just  $f$  being bounded as  $\text{im}(\tau) \rightarrow \infty$ .

### Definition. Modular forms of level $SL_{\mathbb{Z}}(2)$ on upper-half plane

A weakly modular function

#### Definition. Weakly modular forms of level $SL_{\mathbb{Z}}(2)$ on upper-half plane

Let  $k \in \mathbb{N}$ . A meromorphic function

$$f : H \rightarrow \mathbb{C}$$

is **weakly modular** of weight  $k$  if

$$\tau \in \mathbb{H}, \begin{bmatrix} a & b \\ c & d \end{bmatrix} \in SL_{\mathbb{Z}}(2) \implies f\left(\begin{bmatrix} a & b \\ c & d \end{bmatrix}(\tau)\right) = (c\tau + d)^k f(\tau)$$

is a **modular form of weight  $k$**  if  $f$  is holomorphic on  $H$  and at  $\infty$ .

The  $\mathbb{C}$ -vector space of modular forms of wight  $k$  is denoted by

$$\text{Modular}_k(SL_{\mathbb{Z}}(2))$$

and

$$\text{Modular}(SL_{\mathbb{Z}}(2)) := \bigoplus_{k \in \mathbb{Z}} \text{Modular}_k(SL_{\mathbb{Z}}(2))$$

forms a  $\mathbb{Z}$ -graded  $\mathbb{C}$ -algebra.

---

Current note has 0 direct children and 0 total descendants.

- [stamp](#) stamp
  - [Rf](#) subobjects of and functions on  $\mathbb{R}^n, T^n, S^n, \mathbb{C}^n$ 
    - [1Hol](#) Holomorphic functions on spaces over  $\mathbb{C}$  of dimension 1
      - [modular](#) Modular forms

And it has 22 siblings.

- [stamp](#) stamp
  - [Rf](#) subobjects of and functions on  $\mathbb{R}^n, T^n, S^n, \mathbb{C}^n$ 
    - [1Hol](#) Holomorphic functions on spaces over  $\mathbb{C}$  of dimension 1
      - [approx rat](#) Approximation of holomorphic functions on  $\mathbb{C}$  by rational functions
      - [embedding into C3](#) Embedding unit disk  $D$  into  $\mathbb{C}^3$
      - [factorization](#) Factorization of holomorphic functions on  $\mathbb{C}$
      - [global](#) Global holomorphic functions
      - [Hol](#) Equivalent descriptions of holomorphic functions
      - [Mer](#) Meromorphic functions on the complex plane
      - [Mer at infinity](#) Holomorphic functions meromorphic at infinity
      - [modular](#) Modular forms
      - [recons mer from stalk](#) Reconstructing meromorphic functions from stalks
      - [reflection](#) Extending holomorphic functions by reflections
      - [rotation symmetrizer](#) Rotational symmetrization of holomorphic functions
      - [sheaf](#) Sheaf of holomorphic functions on  $\mathbb{C}$
      - [space](#)  $\mathcal{O}(U)$
      - [space C](#)  $\mathcal{O}(\mathbb{C})$
      - [space D](#)  $\mathcal{O}(D)$
      - [space D closed](#)  $\mathcal{O}(\overline{D})$
      - [space D cnt bd](#)  $\mathcal{O}(D) \cap \mathcal{C}(\overline{D})$
      - [space D L2](#)  $\mathcal{O} \cap L^2(D)$
      - [space H](#)  $\mathcal{O}^p(H_{\mathbb{U}}^2)$

- space  $L^p$   $\mathcal{O} \cap L^p$
- space  $S^1$   $\mathcal{O}(S^1)$
- zeros and singularities Zeros and singularities of holomorphic functions